

AD-A037 387

EDGEWOOD ARSENAL ABERDEEN PROVING GROUND MD  
DOSE-RESPONSE EFFECTS OF INTRAVENOUS THIAMINE HYDROCHLORIDE ON --ETC(U)  
MAR 77 J JOSSELSO, F R SIDELL

F/G 6/15  
ON --ETC(U)

UNCLASSIFIED

EB-TR-76117


NL

1 of 1  
ADAO37387



END

DATE  
FILMED  
4-77



ADA 037387

AD.

EDGEWOOD ARSENAL TECHNICAL REPORT  
EB-TR-76117

12  
NW

DOSE-RESPONSE EFFECTS OF INTRAVENOUS THIAMINE HYDROCHLORIDE  
ON PRALIDOXIME PHARMACOKINETICS IN MAN

By

John Josselson, M.D., MAJ, MC, USAR  
Frederick R. Sidell, M.D.

Biomedical Laboratory

March 1977



AD NO. \_\_\_\_\_  
DDC FILE COPY,



DEPARTMENT OF THE ARMY  
Headquarters, Edgewood Arsenal  
Aberdeen Proving Ground, Maryland 21010



Approved for public release; distribution unlimited.

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER EB-TR-76117	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DOSE-RESPONSE EFFECTS OF INTRAVENOUS THIAMINE HYDROCHLORIDE ON PRALIDOXIME PHARMACOKINETICS IN MAN		5. TYPE OF REPORT & PERIOD COVERED Technical Report January 1975 - March 1975
7. AUTHOR(s) John Josselson, M. D., MAJ, MC, USAR Frederick R. Sidell, M.D.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Commander, Edgewood Arsenal Attn: SAREA-BLO Aberdeen Proving Ground, Maryland 21010		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Commander, Edgewood Arsenal Attn: SAREA-TS-R Aberdeen Proving Ground, Maryland 21010		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1W762710AD2502
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 14p		12. REPORT DATE March 1977
		13. NUMBER OF PAGES 19
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited. Copies available from: National Technical Information Service, Springfield, Virginia 22151.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES  Medical defense against chemical agents, prophylaxis and therapy for lethal agents		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Pralidoxime chloride      Plasma concentration Thiamine hydrochloride      Urinary excretion Human subjects		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  Five subjects were given pralidoxime chloride (5 mg/kg, iv) alone or at the end of the first hour of a 3-hour infusion of thiamine hydrochloride (100 mg/hr on one occasion and 200 mg/hr on another). Plasma concentrations of pralidoxime were significantly higher when the oxime was given during a thiamine infusion; plasma concentrations of the oxime were increased slightly more when the higher dose of thiamine was infused. Total oxime excretion was equal in the three groups, but the thiamine-treated groups excreted less oxime during the first 6 hours after drug administration. This suggests that thiamine caused higher concentrations of oxime in the blood by blocking the renal excretion of the drug.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

401 007

7/8



## PREFACE

The work described in this report was authorized under Project/Task 1W762710AD2502, Medical Defense Against Chemical Agents, Prophylaxis and Therapy for Lethal Agents. This work was started in January 1975 and completed in March 1975.

The volunteers in these tests are enlisted US Army personnel. These tests are governed by the principles, policies, and rules for medical volunteers as established in AR 70-25 and the Declaration of Helsinki.

The use of trade names in this report does not constitute an official endorsement or approval of the use of such commercial hardware or software. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, Edgewood Arsenal, Attn: SAREA-TS-R, Aberdeen Proving Ground, Maryland 21010; however, DDC and the National Technical Information Service are authorized to reproduce the document for US Government purposes.

ADDITIONAL FOR	
NTIS	ADDC SECTION <input checked="" type="checkbox"/>
DIC	DDP SECTION <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
IDENTIFICATION	<input type="checkbox"/>
BY: _____	
DISTRIBUTION/AVAILABILITY CODES	
DIC	ADDC SECTION <input type="checkbox"/>
DIC	ADDC SECTION <input type="checkbox"/>
A	

## CONTENTS

	<u>Page</u>
I. INTRODUCTION . . . . .	7
II. METHODS . . . . .	7
A. Subjects . . . . .	7
B. Design . . . . .	7
C. Controls . . . . .	7
D. Thiamine Treatment (100 mg/hr) . . . . .	8
E. Thiamine Treatment (200 mg/hr) . . . . .	8
F. Calculations . . . . .	8
III. RESULTS . . . . .	9
A. Clinical Findings . . . . .	9
B. Plasma Concentration . . . . .	9
C. Urinary Excretion of Pralidoxime . . . . .	9
D. Renal Clearance . . . . .	9
IV. KINETIC DATA . . . . .	12
A. Plasma Clearance . . . . .	12
B. Volumes of Distribution . . . . .	12
C. Plasma Half-Life . . . . .	12
D. Equilibrium Constants . . . . .	13
V. DISCUSSION . . . . .	13
LITERATURE CITED . . . . .	15
DISTRIBUTION LIST . . . . .	17

## DOSE-RESPONSE EFFECTS OF INTRAVENOUS THIAMINE HYDROCHLORIDE ON PRALIDOXIME PHARMACOKINETICS IN MAN

### I. INTRODUCTION.

The cholinesterase reactivator, pralidoxime chloride (N-methylpyridinium-2-aldoxime), remains the drug of choice in this country for the definitive treatment of organophosphate anti-cholinesterase poisoning, and is generally used in conjunction with one of the belladonna alkaloids, usually atropine. The major limitations in the use of all of the currently available oximes have been the production of gastrointestinal side effects (nausea and vomiting with the risk of aspiration in an unconscious patient), hypertension when used in large doses (above 20 mg/kg), and the short half-life and rapid excretion of the drugs. Previous studies in this laboratory have shown that thiamine hydrochloride, U.S.P., when administered either intravenously or intramuscularly immediately before or concomitant with the administration of a rapid intravenous injection of pralidoxime, will significantly increase the plasma half-life and plasma oxime concentration and delay the excretion of the oxime, presumably without altering the mode of excretion.\*,<sup>1</sup> The study presented here was directed toward determining whether or not the response to thiamine was a dose-related event. Thus, subjects were tested according to previous protocols but, on this occasion, using two different doses of thiamine HCl.

### II. METHODS.

#### A. Subjects.

The subjects were US Army enlisted personnel who agreed to the protocol after thorough explanation and discussion. Screening included complete physical and laboratory examinations [chest X-ray, ECG, complete blood count, routine urinalysis, blood urea nitrogen, serum creatinine, liver function tests (SGOT, alkaline phosphatase, serum bilirubin)]. A detailed history of allergy was obtained and intradermal scratch testing with thiamine was performed to exclude potentially hypersensitive subjects.

#### B. Design.

Five subjects participated and received two different doses of thiamine, each serving as his own control. Subjects entered the ward on the evening before the study, ate a light breakfast about 1 hour before the study began, and drank about 1 to 2 liters of fluid in the 2 hours that preceded drug administration.

#### C. Controls.

At the start of testing, a continuous infusion of normal saline into the antecubital vein was begun and continued for 3 hours (200 ml/hr). One hour after the infusion was started, each subject received an intravenous injection of pralidoxime chloride (2-formyl-1-methylpyridinium chloride oxime),\*\* 5 mg/kg, for 2 minutes through the scalp needle of the infusion line.

\* Josselson, J., and Sidell, F. R. EB-TR-76115. The Effect of Thiamine Hydrochloride on Pralidoxime Pharmacokinetics in Man. Submitted for publication.

\*\*Protopam, Ayerst Laboratories, New York, New York. 10017.



D. Thiamine Treatment (100 mg/hr).

At the start of testing, a continuous infusion containing normal saline and thiamine hydrochloride\* was begun; the thiamine was delivered at a rate of 100 mg/hr over a 3-hour period (0.5 mg thiamine/ml of normal saline, 200 ml/hr). One hour after the infusion was begun, each subject received an injection of pralidoxime, 5 mg/kg, as in the control test.

E. Thiamine Treatment (200 mg/hr).

This trial was conducted in the same manner except that the thiamine was delivered at a rate of 200 mg/hr over a 3-hour period (1.0 mg thiamine/ml of normal saline, 200 ml/hr).

The study was randomized and each administration was performed 1 week apart, so that on any given day pralidoxime chloride alone and with both doses of thiamine was administered.

Timed urine collections were obtained from all subjects at 1.5, 3.0, 6.0, and 24 hours. Plasma specimens were obtained through an indwelling scalp needle in a vein in the opposite arm, kept open with heparin, at 0.05, 0.10, 0.15, 0.25, 0.50, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 5.0, and 6.0 hours after the injection of oxime. Plasma and urine were analyzed by the method of Groff, *et al.*<sup>2</sup> for oxime. Subjective responses to both thiamine and pralidoxime were monitored. Subjects were kept in bed during the 3-hour infusion period, thereafter, they were allowed to walk about.

F. Calculations.

The serial plasma oxime concentration, when plotted against time, fit the curve described by the bi-exponential equation 1 which describes the disposition of drug in a two-compartment

$$C_p = Ae^{-\alpha t} + Be^{-\beta t} \quad (1)$$

model. In this equation,  $C_p$  is the plasma concentration,  $t$  is the time,  $A$  and  $B$  represent the zero time intercepts of the linear components of the rapid and slow portions of the plasma decay curve, and  $\alpha$  and  $\beta$  represent the slopes of these two lines divided by -2.303. A more detailed description of the derivations of this equation is provided by Wagner<sup>3</sup> and devised with respect to pralidoxime in studies by Swartz *et al.*<sup>4</sup> and Sidell.<sup>5</sup> The computer program NONLIN<sup>6</sup> was used to estimate the parameters of the equation. From these the following were calculated:

1. The half-life of the rapid equilibrium phase ( $t_{1/2} \alpha$ ) and the slower elimination phase ( $t_{1/2} \beta$ )
2. The volumes of distribution of the central compartment ( $V_1$ ), the peripheral compartment ( $V_2$ ), and the total volume ( $V_D$ )
3. The rate constants for drug movement from central to peripheral compartment ( $K_{12}$ ), from peripheral to central compartment ( $K_{21}$ ), and for elimination from the central compartment ( $K_{13}$ )
4. The plasma and renal clearance of pralidoxime. The plasma clearance was

\*Thiamine Hydrochloride Injection U.S.P., 10 mg/ml, Natcon Chemical Co.

calculated by dividing the dose of pralidoxime by the area under the plasma concentration versus time curve (which was obtained from the integral of equation 1), and the renal clearance was obtained by dividing the amount of pralidoxime excreted into the urine during a given time period by the area under the curve for that time interval.

### III. RESULTS.

#### A. Clinical Findings.

Both thiamine and pralidoxime were well tolerated by all subjects with no major untoward side effects. No changes in vital signs were observed during the course of all three trials. All five subjects, during the control testing, complained of subjective visual disturbances, ranging from "blurring" to "heavy eyelids;" one complained of actual diplopia. These symptoms occurred only transiently in controls (lasting up to 10 minutes), were present in four of the five subjects during treatment with the low dose of thiamine (100 mg/hr), lasting up to 1 hour, and were present in all subjects receiving 200 mg/hr of thiamine, persisting up to 2 hours in four of the five subjects. Three subjects complained of prolonged lethargy and drowsiness following the high dose of thiamine, but these symptoms disappeared by the morning after testing.

#### B. Plasma Concentration.

The mean plasma concentrations of pralidoxime are plotted in the following figure. After 3 minutes (0.05 hr), statistically significant differences were noted when controls were compared with both the low-dose ( $T_1$ ) thiamine group ( $p < 0.05$ , paired t test) and the high-dose ( $T_2$ ) thiamine group ( $p < 0.05$ , paired t test), and when the  $T_1$  group was compared to the  $T_2$  group ( $p < 0.05$ , paired t test) up to hour 5 of plasma collection.

#### C. Urinary Excretion of Pralidoxime (Table 1).

Over the 24-hour period of urine collection, there was no significant difference in total oxime excretion among the three groups. The controls (C) excreted 74.8% ( $\pm 12$  SD); the low dose thiamine group ( $T_1$ ) 66.7% ( $\pm 6.7$  SD); and the high-dose thiamine group ( $T_2$ ) 69.1% ( $\pm 7.6$  SD) of their doses of pralidoxime. However, over the first 6 hours, the thiamine-treated groups eliminated significantly less oxime [C 71% ( $\pm 11$  SD);  $T_1$  49.9% ( $\pm 8.7$  SD);  $T_2$  43.6% ( $\pm 6.7$  SD)] when compared to controls ( $p < 0.01$ , paired t test), but not when compared to each other. Additionally, over the last 18 hours (6 to 24 hours) of timed urine collection, significantly more oxime appeared in the urine of both thiamine-treated groups when compared to controls and to each other [C 3.4% ( $\pm 3$  SD);  $T_1$  16.8% ( $\pm 5.3$  SD);  $T_2$  25.5% ( $\pm 7$  SD) (C vs  $T_1$   $p < 0.01$ ; C vs  $T_2$   $p < 0.01$ ;  $T_1$  vs  $T_2$   $p < 0.05$ ; paired t test)].

#### D. Renal Clearance.

Mean renal clearance data are presented in table 1. The 24-hour renal clearance of oxime was reduced significantly in both thiamine-treated groups, when compared to controls and to each other (C vs  $T_1$ ,  $p < 0.05$ ; C vs  $T_2$ ,  $p < 0.05$ ;  $T_1$  vs  $T_2$ ,  $p < 0.05$ , paired t test).

Table 1. Urinary Recovery of Pralidoxime

Time hr	C	T <sub>1</sub>	T <sub>2</sub>	p <sup>a</sup>
A. Mean Cumulative Excretion of Pralidoxime Over 24 Hours <sup>b</sup>				
0-1.5	77.4± <sup>c</sup>	45.3±	34.6±	0.01
	5.5	5.8	3.5	0.001
				0.05
0-3	0.5±	54.6±	48.9±	0.001
	4.8	7.0	6.1	0.001
				NS
0-6	95.1±	74.5±	63.4±	0.01
	4.9	8.9	8.4	0.01
				0.05
0-24	100	100	100	-
B. Mean Urinary Excretion of Pralidoxime for Each Collection Period <sup>b</sup>				
0-1.5	77.4±	43.5±	34.6±	0.01
	5.5	5.8	3.5	0.001
				0.05
0-5.3	13.1±	11.1±	14.3±	NS
	3.3	1.9	2.7	NS
				NS
3-6	5.0±	19.8±	14.5±	0.001
	3.6	3.4	4.6	0.02
				NS
6-24	4.5±	25.5±	36.8±	0.01
	4.3	8.9	8.7	0.01
				0.05
Total	100	100	100	-

<sup>a</sup> C versus T<sub>1</sub>.C versus T<sub>2</sub>.T<sub>1</sub> versus T<sub>2</sub>.<sup>b</sup> Expressed as percent of the total amount of oxime excreted in the urine.<sup>c</sup> Mean ± SD.

NOTE: NS - not sufficient.




Figure. A Plot of the Mean Concentrations of Pralidoxime



#### IV. KINETIC DATA.

##### A. Plasma Clearance.

Similarly, the plasma clearance of pralidoxime was also significantly reduced by both levels of thiamine administration (table 2: C vs T<sub>1</sub>,  $p < 0.05$ ; C vs T<sub>2</sub>,  $p < 0.02$ ; T<sub>1</sub> vs T<sub>2</sub>,  $p < 0.05$ , paired t test).

Table 2. Mean Renal and Plasma Clearance Values of Pralidoxime Chloride in Five Men

Renal clearance (ml/min)	603 ± 352*	245 ± 83.7	178 ± 55.1	<0.05	<0.05	<0.05
Plasma clearance (ml/min)	774 ± 308	364 ± 97.8	258 ± 76.5	<0.05	<0.02	<0.05
Area under curve (mcg. hr/ml)	7.04 ± 1.85	14.36 ± 2.86	20.93 ± 6.89	<0.001	<0.001	NS

\*Mean ± SD.

##### B. Volumes of Distribution.

The remaining pharmacokinetic data are presented in table 3. Following administration of the high dose of thiamine, there was a significant decrease in the total volume of distribution when compared to controls and to the low dose (C vs T<sub>2</sub>,  $p < 0.02$ ; T<sub>1</sub> vs T<sub>2</sub>,  $p < 0.01$ , paired t test). These results were characterized by a large decrease in the peripheral (extravascular) compartment (V<sub>2</sub>), (C vs T<sub>2</sub>,  $p < 0.02$ ; T<sub>1</sub> vs T<sub>2</sub>,  $p < 0.02$ , paired t test), and essentially no change in the central compartment (V<sub>1</sub>). Though the data conflict somewhat with those previously reported\* in which low dose also decreased the total and peripheral volumes of distribution while slightly increasing the central volume, nevertheless, the trends are distinctly similar and emphasize the additional effects of the higher dose of thiamine (200 mg/hr).

##### C. Plasma Half-Life (Table 2).

Following thiamine administration, significant increases in the long half-life of elimination ( $t_{1/2\beta}$ ) occurred in both thiamine-treated groups when compared to controls, but not to each other (C vs T<sub>1</sub>,  $p < 0.05$ ; C vs T<sub>2</sub>,  $p < 0.05$ ; T<sub>1</sub> vs T<sub>2</sub>, NS). Our data substantiate our previous findings\* that thiamine hydrochloride (100 mg/hr) did prolong the short (or equilibration phase) half-time ( $t_{1/2\alpha}$ ), but data comparing the control dose with the high dose of thiamine and comparing the high and low doses of thiamine showed no significance.

\*Josselson, J., and Sidell, R. R. EB-TR-76115. The Effects of Thiamine Hydrochloride on Pralidoxime Pharmacokinetics in Man. Submitted for publication.



Table 3. Pharmacokinetic Parameters

	Control	Thiamine 1	Thiamine 2	Statistical significance (p)		
				C vs T <sub>1</sub>	C vs T <sub>2</sub>	T <sub>1</sub> vs T <sub>2</sub>
"Short" Half-Time (t <sub>1/2</sub> ) (hr)	0.07 ± 0.005*	0.14 ± 0.03	0.14 ± 0.11	<0.01	NS	NS
"Long" Half-Time (t <sub>1/2</sub> ) (hr)	0.99 ± 0.13	1.46 ± 0.24	1.52 ± 0.47	<0.05	<0.05	NS
k <sub>21</sub> (hr <sup>-1</sup> )	2.37 ± 0.44	1.94 ± 0.43	2.64 ± 1.40	NS	NS	NS
k <sub>13</sub> (hr <sup>-1</sup> )	2.94 ± 0.75	1.32 ± 0.26	1.21 ± 0.47	<0.01	<0.01	NS
k <sub>12</sub> (hr <sup>-1</sup> )	4.95 ± 0.40	2.49 ± 0.78	3.21 ± 1.85	<0.001	NS	NS
V <sub>1</sub> (central) (ml/kg)	259 ± 40.7	275 ± 39.5	232 ± 71.4	NS	NS	NS
V <sub>2</sub> (peripheral) (ml/kg)	563 ± 178	335 ± 29.5	257 ± 101	NS	<0.02	<0.02
V <sub>D</sub> (total) (ml/kg)	822 ± 211	618 ± 48.8	469 ± 49.1	NS	<0.02	<0.01

\*Mean ± SD.

D. Equilibrium Constants (Table 3).

Following the low dose of thiamine, a highly significant decrease occurred in the rate constant (K<sub>12</sub>) governing movement of oxime from the central to the peripheral compartments (C vs T<sub>1</sub>, p < 0.001, paired t test), but no trends were identified when the high dose of thiamine was compared to the low dose or controls (C vs T<sub>2</sub>, p: NS; T<sub>1</sub> vs T<sub>2</sub>, p: NS, paired t test)

Significant changes also occurred in the rate constant (K<sub>13</sub>) when controls were compared to both thiamine levels (C vs T<sub>1</sub>, p < 0.01; C vs T<sub>2</sub>, p < 0.01, paired t test), but no significant differences were noted between the two thiamine-treated groups. The changes noted reflected the decrease in renal (and plasma) clearance before and after pretreatment with thiamine.

V. DISCUSSION.

That the half-life, renal and plasma clearances, and pharmacokinetics can be altered by prior or simultaneous administration of thiamine hydrochloride intramuscularly or intravenously has been demonstrated conclusively by this and previous studies.\*<sup>1</sup> The data presented here are in close agreement with our previous report, and the minor differences [especially with respect to half-life (t<sub>1/2β</sub>), volumes of distribution, and rate constants] appear to be due to the limited size of our population (n=5). Nevertheless, the experience in this laboratory over the past 2 years with a total of 20 subjects substantiates our findings.\*<sup>1</sup>

\*Josselson, J., and Sidell, F. R. EB-TR-76115. The Effects of Thiamine Hydrochloride on Pralidoxime Pharmacokinetics in Man. Submitted for publication.

Likewise, the fact that the higher dose of thiamine hydrochloride (200 mg/hr) should be more effective than the lower dose (100 mg/hr) was also not unexpected. Though the higher dose did not significantly alter the long plasma half-life ( $t_{1/2}$ ), it did increase it slightly. More importantly, however, the higher dose did significantly decrease both renal and plasma clearance of oxime when compared to the lower dose and, perhaps even more importantly, did maintain significantly higher plasma levels of oxime over the first 4 hours of treatment.

Though pralidoxime is well recognized as an effective cholinesterase reactivator in the treatment of organophosphate nerve agent and insecticide poisoning, few, if any, objective criteria have been established to indicate what parameters one can monitor to achieve successful therapy. Clearly, the patient must be treated rapidly (within at least 2 hours). Available information on antidotes suggests that the optimal plasma concentration for effective regeneration of the inhibited enzyme is approximately 4 mg%. If this is the case, then the intravenous administration of the high dose of thiamine (200 mg/hr infusion) appears to be advantageous over low-dose therapy (100 mg/hr) and will maintain levels of 4 mg% for up to 1 hour. Low-dose therapy holds that level for only 30 minutes, and when there is no pretreatment, the plasma concentration falls below the 4 mcg/ml level in 15 minutes (at oxime doses of 5 mg/hr). At this point, oxime re-administration may be necessary.

While the higher dose of thiamine did markedly prolong the clinical side effects of oxime and did produce more lethargy than the low-dose therapy, it was, in general, excellently tolerated by all subjects. Use of this procedure may avoid the problems involved with larger doses of oxime (i.e., hypertension and gastrointestinal disturbances).

Studies are now under way to determine whether drip infusion of pralidoxime and thiamine hydrochloride over 1 to 2 hours will be statistically effective. Once studies have been performed in an adequately simulated casualty situation, better determination of the effectiveness of thiamine hydrochloride should be possible.

### LITERATURE CITED

1. Swartz, R. D., and Sidell, F. R. Renal Tubular Secretion of Pralidoxime in Man. *Proc. Soc. Exper. Biol. Med.*, *146*, 419-424 (1974).
2. Groff, W. A., and Ellin, R. I. A New and Rapid Determination of Pyridinium Aldoximes in Blood and Urine. *Clin. Chem.* *15*, 72-83 (1969).
3. Wagner, J. C. Pharmacokinetics, Grosse Pointe, Michigan, 1969, J. M. Richard Laboratories.
4. Swartz, R. D., and Sidell, F. R. Effects of Heat and Exercise on the Elimination of Pralidoxime in Man. *Clin. Pharmacol. Therap.* *14*, 83-89 (1973).
5. Sidell, F. R., *et al.* Toxogonin and Pralidoxime: Kinetic Comparison After Intravenous Administration to Man. *J. Pharmaceut. Sci.* *61*, 1765-1769 (1972).
6. Metzler, C. M. A Users Manual for NONLIN, The Upjohn Company, November 25, 1969.

## DISTRIBUTION LIST 5

Names	Copies	Names	Copies
EDGEWOOD ARSENAL		Director	
FOREIGN INTELLIGENCE OFFICER	1	Defense Civil Preparedness Agency	
CHIEF, LEGAL OFFICE	1	Attn: RE	1
CHIEF, SAFETY OFFICE	1	Attn: PO(DC)	1
CHIEF, PLANS OFFICE		Washington, DC 20301	
Attn: SAREA-PL	1	CINCUSAREUR	
CDR, US ARMY TECHNICAL ESCORT CENTER	1	Attn: AEAGC-DS	1
AUTHOR'S COPY, Biomedical Laboratory	2	APO New York 09403	
BIOMEDICAL LABORATORY		Chief, Office of Research,	
Attn: SAREA-BL	1	Development & Acquisition	
Attn: SAREA-BL-M	7	Attn: DAMA-CSM-CM	1
Attn: SAREA-BL-Q	1	Attn: DAMA-ARZ-D	1
Attn: SAREA-BL-V	1	Washington, DC 20310	
CHEMICAL LABORATORY		Headquarters, Sixth US Army	
Attn: SAREA-CL	1	NBC Office	
Attn: SAREA-CL-P	1	Attn: AFKC-OP-PN	1
Attn: SAREA-CL-TE	1	Presidio of San Francisco, CA 94129	
DEVELOPMENT AND ENGINEERING DIRECTORATE		CINCUSAREUR	
Attn: SAREA-DE-D	1	Attn: AEAMD-PM	1
Attn: SAREA-DE-DB	1	APO New York 09403	
Attn: SAREA-DE-EP	1	OFFICE OF THE SURGEON GENERAL	
MANUFACTURING TECHNOLOGY DIRECTORATE		HQDA (SGRD-HR)	1
Attn: SAREA-MT-D	1	Forrestal Building	
Attn: SAREA-MT-P	1	WASH DC 20314	
DIRECTOR OF PRODUCT ASSURANCE		US ARMY HEALTH SERVICE COMMAND	
Attn: SAREA-PA-A	1	Commander	
Attn: SAREA-PA-P	1	US Army Environmental Hygiene Agency	
Attn: SAREA-PA-Q	1	Attn: HSE-AA-L, Librarian, Bldg 2100	1
DIRECTOR OF TECHNICAL SUPPORT		APG-Edgewood Area	
Attn: SAREA-TS-R	2	Superintendent	
Attn: SAREA-TS-L	3	Academy of Health Sciences	
Attn: SAREA-TS-E	1	US Army	
DEPARTMENT OF DEFENSE		Attn: HSA-CDC	1
Administrator		Attn: HSA-IHE	1
Defense Documentation Center		Fort Sam Houston, TX 78234	
Attn: Accessions Division	12	US ARMY MATERIEL DEVELOPMENT AND	
Cameron Station		READINESS COMMAND	
Alexandria, VA 22314		Commander	
Director		US Army Materiel Development and	
Defense Intelligence Agency		Readiness Command	
Attn: DB-4G1	1	Attn: DRCDE-DM	1
Washington, DC 20301		Attn: DRCLDC	1
DEPARTMENT OF THE ARMY		5001 Eisenhower Ave	
HQDA (DAMO-SSC)	1	Alexandria, VA 22333	
WASH DC 20310			



# DISTRIBUTION LIST 5 (Contd)

Names	Copies	Names	Copies
Commander US Army Foreign Science & Technology Center Attn: DRXST-ISI 220 Seventh St., NE Charlottesville, VA 22901	2	Headquarters US Army Military Police School/Training Center Attn: ATSJ-TD-CR Fort McClellan, AL 36205	1
Commander DARCOM, STITEUR Attn: DRXST-ST1 Box 48, APO New York 09710	1	Commander US Army Infantry Center Attn: ATZB-CD-MS-C Fort Benning, GA 31905	1
Commander US Army Science & Technology Center-Far East Office APO San Francisco 96328	1	Commandant US Army Ordnance Center & School Attn: ATSL-CD-MS Attn: ATSL-DTG APG-Aberdeen Area	1 1
Redstone Scientific Information Center Attn: Chief, Documents US Army Missile Command Redstone Arsenal, AL 35809	1	US ARMY TEST & EVALUATION COMMAND Record Copy CDR, APG Attn: STEAP-AD-R/RHA APG-Edgewood Area, Bldg E5179	1
Director DARCOM Field Safety Activity Attn: DRXOS-C Charlestown, IN 47111	1	CDR, APG Attn: STEAP-TL APG-Aberdeen Area	1
US ARMY ARMAMENT COMMAND		Commander US Army Test & Evaluation Command Attn: DRSTE-FA APG-Aberdeen Area	1
Commander US Army Armament Command Attn: DRSAR-ASN Attn: DRSAR-IMBC Rock Island, IL 61201	1 1	Commander US Army Tropic Test Center Attn: STETC-MO-A (Tech Library) APO New York 09827	1
Commander US Army Dugway Proving Ground Attn: Technical Library, Docu Sect Dugway, UT 84022	1	Commander Dugway Proving Ground Attn: STEDP-PO Dugway, UT 84022	1
Commander Rocky Mountain Arsenal Attn: SARRM-EA Denver, CO 80240	1	DEPARTMENT OF THE NAVY Chief of Naval Research Attn: Code 443 800 N. Quincy Street Arlington, VA 22217	1
US ARMY TRAINING & DOCTRINE COMMAND		Commander Naval Facilities Engineering Command Attn: Code 03 200 Stovall Street Alexandria, VA 22332	1
Commandant US Army Infantry School Combat Support & Maintenance Dept. Attn: NBC Division Fort Benning, GA 31905	1	Commander Naval Explosive Ordnance Disposal Facility Attn: Army Chemical Office Indian Head, MD 20640	1
Commandant US Army Missile & Munitions Center & School Attn: ATSK-CD-MD Attn: ATSK-DT-MU-EOD Redstone Arsenal, AL 35809	1 1		



# DISTRIBUTION LIST 5 (Contd)

Names	Copies	Names	Copies
Commander Naval Surface Weapons Center Attn: Tech Lib & Info Svcs Br White Oak Laboratory Silver Spring, MD 20910	1	HQ Foreign Technology Division (AFSC) Attn: PDRR Wright-Patterson AFB, OH 45433	1
Commander Naval Surface Weapons Center Dahlgren Laboratory Attn: DX-21 Dahlgren, VA 22448	1	Commander Aeronautical Systems Division Attn: ASD/AELD Wright-Patterson AFB, OH 45433	1
Commander Nuclear Weapons Training Group, Atlantic Naval Air Station Attn: Code 21 Norfolk, VA 23511	1	HQ, USAF/SGPR Forrestal Bldg WASH DC 20314	1
Chief, Bureau of Medicine & Surgery Department of the Navy Attn: CODE 5 Washington, DC 20372	1	HQ, USAF/RDPN WASH DC 20330	1
DEPARTMENT OF THE AIR FORCE		HQ, Ogden ALC/MMWRA Hill AFB, UT 84406	1
Air University Library Attn: AUL/LSE-8879 Maxwell AFB, AL 36112	1	Director Air Force Inspection and Safety Center Attn: IGD(AFISC/SEV) Norton AFB, CA 92409	1
		Commander Armament Development & Test Center Attn: DLOSL (Technical Library) Eglin AFB, FL 32542	1
		NORAD Combat Operations Center/DBN Cheyenne Mtn Complex, CO 80914	1